

Gamma-ray Pulsars in a Modified Polar Cap Scenario

B. Rudak and J. Dyks

N. Copernicus Astronomical Center, Rabiańska 8, 87-100 Toruń, Poland

Abstract. We present a polar-cap model which incorporates a likely acceleration of Sturrock pairs with their subsequent contribution to gamma-ray luminosity L_γ . This model reproduces L_γ for seven pulsars detected with *Compton Gamma Ray Observatory* experiments, avoiding at the same time the problem of the empirical gamma-ray death line of Arons (1996).

Also, we estimate the efficiency of reversing newly created positrons by residual longitudinal electric field. Over the wide range of spin-down luminosity values the predicted polar-cap X-ray luminosity $L_X(pc)$ goes as $\propto L_{sd}^{0.6}$. Model calculations for B0823+26, B0950+08, B1929+10, and J0437-4715 are compared with existing observational constraints on thermal X-ray components.

1. Introduction

We present a semi-empirical polar cap model which reproduces the gamma-ray luminosities inferred for seven pulsars (we dub them *Seven Samurai*) detected with the CGRO experiments: Crab, B1509-58, Vela, B1951+32, B1706-44, Geminga, and B1055-52. The model avoids the problem of the empirical gamma-ray death line of Arons (1996). We also address the question of expected thermal X-ray emission due to polar cap reheating by reversed particles. With numerical calculations used to obtain L_γ we tried to reproduce recent *ROSAT* and *ASCA* results.

2. Gamma-ray Luminosities

The model (see Rudak & Dyks 1997 for details) is based on polar cap activity triggered by ultrarelativistic primary electrons as described by Daugherty & Harding (1982). Our two basic assumptions are:

1) The energy of primary electrons E_0 is only a few times higher than the threshold energy required to induce pair creation in a dipolar magnetic field of surface strength B_{12} Teragauss, with other restrictions applied when necessary:

$$E_0 = \min\{\zeta \cdot E_{\min}, E_W, E_{\max}\}, \quad (1)$$

where ζ satisfies $2 \lesssim \zeta \lesssim 5$, and $E_{\min} = 1.2 \times 10^7 B_{12}^{-1/3} P^{1/3}$, $E_W = 1.2 \times 10^7 B_{12} P^{-2}$, $E_{\max} = 4.6 \times 10^7 B_{12}^{1/4} P^{-1/8}$ (Rudak & Ritter 1995).




Figure 1. Total L_γ inferred from *EGRET*, *COMPTEL*, and *OSSE* is shown for seven pulsars (filled dots). Open triangles are the *EGRET* upper limits (Nel et al.1996). Filled triangle marks the *EGRET* U.L. for the millisecond pulsar J0437-4715. The dashed line represents $L_\gamma = 10^{16} \cdot L_{sd}^{1/2}$. The upper solid line is the evolutionary track calculated for $B = 10^{12}$ G, the lower one is for 10^9 G.

2) Out of $n_\pm(E_0)$ Sturrock pairs produced per each primary electron a fraction η_γ is subject to further acceleration and contributes to pulsar's gamma-ray emission.

We obtain thus a following prescription for L_γ :

$$L_\gamma = \eta_\gamma \cdot E_\pm \cdot n_\pm \cdot \dot{N}_{GJ}, \quad (2)$$

where $E_\pm$ – a characteristic energy attained by secondary particles due to acceleration – is assumed to be of the order of E_0 ; the parameter η_γ will be determined by comparing results of numerical simulations of creation of Sturrock pairs with the *CGRO* results; $\dot{N}_{GJ}$ is the Goldreich-Julian rate of outflow of primary electrons.

The model was confronted with the gamma-ray luminosities for seven pulsars inferred from available data from *OSSE* (Schroeder et al.1994), *COMPTEL* (Carramiñana et al.1995), and *EGRET* (Fierro 1995, Nel et al.1996) experiments (assuming gamma-ray beaming angle $\Omega_\gamma = 1$ sr in all cases). Two evolutionary tracks in the $L_\gamma - L_{sd}$ space calculated for a classical pulsar ($B = 10^{12}$ G), and for a millisecond pulsar ($B = 10^9$ G) are shown in Fig.1.




Figure 2. Thermal X-ray components calculated for *Seven Samurai* along with 4 other classical pulsars (filled dots), and 2 millisecond pulsars (filled triangles). The open triangle is the upper limit for J0437-4715 (Halpern et al.1997). Open circles are possible thermal components for B0823+26 (Becker 1995), B0950+08 and B1929+10 (Wang & Halpern 1997).

For a fixed value of L_{sd} , the predicted L_γ depends rather weakly on magnetic field strength B as long as $10^{11}\text{G} \lesssim B \lesssim 10^{13}\text{G}$, in contrast to other existing models. Due to this property the model reproduces the seven detections equally well as e.g. a simple empirical relation $L_\gamma \propto L_{sd}^{1/2}$ does. At the domain of millisecond pulsars ($\sim 10^8 - 10^9\text{G}$), L_γ drops significantly. The value of parameter η_γ in Eq.2 is $\sim 4 \times 10^{-3}$.

The upper limit for J0437-4715 from *EGRET* is about one order of magnitude above the expected level. Out of four classical pulsars with *EGRET* upper limits visibly below our predictions (see Fig.1) B1046-58 and B1929+10 pose no threat to the model due to existing upper limits set by *COMPTEL*. There is no detailed information available about the other two - B0950+08 and B0656+14.

3. Reheating of the Polar Cap

Some positrons, before reaching a region of further acceleration, may be stopped by local residual longitudinal electric field and directed back to the surface.

Each returning positron acquires on its way an energy comparable to the energy of the primary electron, $E_+ \simeq E_0$, and then deposits it onto the polar cap surface. A heat conductivity inwards the star, and across magnetic lines is negligible and dissipated energy is radiated out on the spot, predominantly in form of thermal X-rays, with the luminosity $L_X(\text{pc})$.

If out of all positrons ($\dot{N}_+ = n_{\pm} \cdot \dot{N}_{\text{GJ}}$) a fraction $\eta_{\downarrow}$ will be reversed, then

$$L_X(\text{pc}) = \eta_{\downarrow} \cdot E_0 \cdot n_{\pm} \cdot \dot{N}_{\text{GJ}}. \quad (3)$$

Suppose now that $\eta_{\downarrow} \propto \epsilon_{\parallel}^{-1}$, where $\epsilon_{\parallel}$ is a mean energy of freshly created pairs retained in their longitudinal motion (Rudak 1997, in preparation).

Fig.2 shows $L_X(\text{pc})$ as a function of spin-down luminosity L_{sd} calculated for *Seven Samurai*, 4 other classical pulsars, and 2 millisecond pulsars. Over the wide range of L_{sd} values the calculated polar-cap X-ray luminosity behaves as $L_X(\text{pc}) \propto L_{\text{sd}}^{0.6}$. This formula is significantly flatter than the empirical relation $L_X \approx 0.001 L_{\text{sd}}$ of Becker & Trümper (1997). Even though some models are able to reproduce this relation with non-thermal X-ray spectra quite satisfactorily (Cheng et al.1997), it seems very likely that actual X-ray spectra consist of two components: a non-thermal one - due to synchrotron radiation, and a thermal one - due to polar-cap reheating. For high L_{sd} pulsars the former component dominates, while at the low- L_{sd} end the latter component becomes visible.

The values of parameter $\eta_{\downarrow}$ from Eq.3 are on average $\sim 10^{-5}$ and $\sim 10^{-3}$ for classical and millisecond pulsars, respectively.

Acknowledgments. This research was financed by the KBN grant 2P03D 00911, and the Jumelage program “Astronomie Pologne”. BR is grateful to Jean Pierre Lasota for his warm hospitality in DARC (Meudon).

References

- Arons, J. 1996, A&AS, 120, 49
- Becker, W. 1995, Ph.D. Thesis (LMU Munich)
- Becker, W., Trümper, J. 1997, A&A, in the press
- Carramiñana A., Bennett K., Buccherri R., et al. 1995 A&A, 304, 258
- Cheng, K.S., Gil, J., Zhang, L. 1997, ApJ, submitted
- Daugherty, J.K., Harding, A.K. 1982, ApJ, 252, 337
- Fierro, J.M. 1995, Ph.D. Thesis (Stanford University)
- Halpern, J.P., Martin, C., Marshall, H.L. 1997, ApJ, 462, 908
- Harding, A.K. 1981, ApJ, 245, 267
- Nel H.I., Arzoumanian Z., Brazier K.T.S., et al. 1996, ApJ, 465, 898
- Rudak, B., Ritter, H. 1994, MNRAS, 267, 513
- Rudak, B., Dyks, J. 1997, MNRAS, submitted
- Schroeder, P.C., Ulmer, M.P., Matz, S.M., et al. 1994, ApJ, 450, 784
- Wang, F.Y.-H., Halpern, J.P. 1997, ApJ, 482, L159